



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A CONDITION THAT A FUNCTION IN A PROJECTIVE SPACE BE RATIONAL*

BY

WILLIAM F. OSGOOD

HURWITZ † has shown that a function of n complex variables which is in general analytic and has no other than unessential singularities is a rational function of the n arguments. The infinite region is that of the theory of functions. A point (z_1, \dots, z_n) lies at infinity if at least one of its coördinates corresponds to the point $z_k = \infty$ in the complex z_k -plane; and such a point (z_1, \dots, z_n) is projected into the finite region by subjecting the individual coördinate or coördinates which become infinite to a linear transformation:

$$Z_k = \frac{1}{z_k},$$

or, more generally,

$$Z_k = \frac{\alpha_k z_k + \beta_k}{\gamma_k z_k + \delta_k}, \quad \alpha_k \delta_k - \beta_k \gamma_k \neq 0, \quad \gamma_k \neq 0.$$

Thus the points of the space of the theory of functions can be represented by n pairs of binary homogeneous variables:

$$[(\zeta_1, \zeta'_1), (\zeta_2, \zeta'_2), \dots, (\zeta_n, \zeta'_n)],$$

the transformations being n independent binary linear transformations:

$$\left. \begin{aligned} \mathbf{Z}_k &= \alpha_k \zeta_k + \beta_k \zeta'_k \\ \mathbf{Z}'_k &= \gamma_k \zeta_k + \delta_k \zeta'_k \end{aligned} \right\} \quad \begin{vmatrix} \alpha_k & \beta_k \\ \gamma_k & \delta_k \end{vmatrix} \neq 0, \quad k = 1, 2, \dots, n.$$

A function is said to be analytic in a point at infinity if, on making the above transformation for as many of the coördinates separately as become infinite, the transformed function is analytic in the new variables at the corresponding point, save as for a removable singularity (hebbare Unstetigkeit). A function

* Presented to the Society February 24, 1912.

† A. HURWITZ, *Journal für Mathematik*, vol. 95 (1883) p. 201. The theorem was stated without proof by WEIERSTRASS.

has an unessential singularity at a point (finite or at infinity), if in the neighborhood of this point, while not remaining finite, it can be expressed as the quotient of two functions, each of which is analytic in the point in question.

§ 1.

In the present paper it is shown that a theorem similar to the above holds for a complex n -fold projective space. Before stating the theorem in precise form we will consider some preliminary definitions and conceptions.

Let a point of the projective space in question be represented by the complex, homogeneous coördinates x_0, x_1, \dots, x_n . These coördinates never vanish simultaneously, and infinite values of the coördinates never enter into consideration. Let $F(x_0, x_1, \dots, x_n)$ be a function single-valued and analytic throughout a complex $(n+1)$ -dimensional continuum \mathfrak{R}_1 lying in the above complex $(n+1)$ -dimensional space \mathfrak{R} of the variables (x_0, x_1, \dots, x_n) , and let F be homogeneous of dimension 0; i. e., if (x_0, x_1, \dots, x_n) is a point of \mathfrak{R}_1 , then all the points $(\lambda x_0, \lambda x_1, \dots, \lambda x_n)$, where λ is any complex number except 0, shall also lie in \mathfrak{R}_1 , and furthermore

$$F(x_0, x_1, \dots, x_n) = F(\lambda x_0, \lambda x_1, \dots, \lambda x_n).$$

Let (a_0, a_1, \dots, a_n) be a point of \mathfrak{R}_1 . Then one of the a 's (a_k , let us say) is different from 0, while on the other hand F is analytic throughout a certain neighborhood of this point:

$$(1) \quad |x_l - a_l| < h, \quad l = 0, 1, 2, \dots, n, \quad 0 < h < |a_k|.$$

If we set $\lambda = x_k^{-1}$, then for the above neighborhood

$$F(x_0, x_1, \dots, x_n) = F\left(\frac{x_0}{x_k}, \frac{x_1}{x_k}, \dots, \frac{x_{k-1}}{x_k}, 1, \frac{x_{k+1}}{x_k}, \dots, \frac{x_n}{x_k}\right).$$

For such values of

$$w_l = \frac{x_l}{x_k}, \quad l = 0, 1, \dots, k-1, k+1, \dots, n,$$

as correspond to the points (x_0, x_1, \dots, x_n) of the neighborhood (1),

$$F(w_0, w_1, \dots, w_{k-1}, 1, w_{k+1}, \dots, w_n)$$

is seen to be an analytic function of the n arguments w .

Finally, let (a_0, a_1, \dots, a_n) be a point of \mathfrak{R} , in which F is not defined. Again, let $a_k \neq 0$, and consider the neighborhood (1), h being suitably restricted. Then F shall be defined in general in the points of this region, and shall be equal to the quotient

$$F(x_0, \dots, x_n) = \frac{Q(x_0, \dots, x_n)}{P(x_0, \dots, x_n)},$$

where ρ is an arbitrary complex number distinct from 0. Thus to every point of S corresponds one point and only one of R , though an infinite number of points in \mathfrak{R} present themselves. If (x_0, x_1, \dots, x_n) is one of the latter points, then the others are $(\lambda x_0, \lambda x_1, \dots, \lambda x_n)$, where $\lambda \neq 0$ is any complex number.

The inverse of (A) is easily written down. Consider the equation in

$$z = \zeta / \zeta' :$$

$$(B_1) \quad x_0 - x_1 z + x_2 z^2 - \dots \pm x_n z^n = 0,$$

or in homogeneous form:

$$(B_2) \quad x_0 \zeta'^n - x_1 \zeta'^{n-1} \zeta + x_2 \zeta'^{n-2} \zeta^2 - \dots \pm x_n \zeta^n = 0.$$

(x_0, x_1, \dots, x_n) being a point of \mathfrak{R} , not all the x 's can vanish, and hence the equation (B_2) determines precisely n (distinct or coincident) pairs of values, — or more properly the ratios of the letters in each pair:

$$(\zeta_1, \zeta'_1), (\zeta_2, \zeta'_2), \dots, (\zeta_n, \zeta'_n),$$

and these in turn, when arranged in all $n!$ possible ways, yield $n!$ points (z_1, z_2, \dots, z_n) in the space S .

We have, then, in (A) and (B) a $(1, n!)$ transformation of S on R , and an $(\infty, n!)$ transformation of S on \mathfrak{R} .

On applying this transformation to the function $F(x_0, \dots, x_n)$, the latter is carried over into a function

$$\Phi(z_1, z_2, \dots, z_n),$$

single-valued at all points of S where it is defined. Moreover, Φ is, as we will presently show, analytic in S where it is defined, and it has no other than unessential singularities in S . It is, therefore, by the theorem cited at the beginning of this paper, a rational function of z_1, \dots, z_n . It is, furthermore, symmetric in these variables, and consequently F is rational in x_0, x_1, \dots, x_n .

To complete the proof, consider first a point $(a) = (a_0, a_1, \dots, a_n)$ of \mathfrak{R} in which $a_n \neq 0$. Let $(b) = (b_1, \dots, b_n)$, where $z_i = b_i$, be one of the points of S into which (a) is carried by the transformation (B_1) . This point will lie in the finite part of S . To every point (x) of a certain neighborhood of (a) , and hence to every point $(w) = (w_0, w_1, \dots, w_{n-1})$ of a certain neighborhood of $(w) = (c) = (c_0, c_1, \dots, c_{n-1})$, where $c_i = a_i/a_n$, will correspond one or more points (z) of a certain neighborhood of (b) ; and conversely, to each point (z) of a neighborhood of (b) will correspond one point (w) of a neighborhood of (c) . Finally, for all pairs of corresponding points we shall have

